

CORRELATION LENGTH OF X-RAY BRIGHTEST ABELL CLUSTERS

MARIO G. ABADI,² DIEGO G. LAMBAS¹ AND HERNÁN MURIEL¹

Grupo de Investigaciones en Astronomía Teórica y Experimental (IATE)
 Observatorio Astronómico de Córdoba, Laprida 854, 5000, Córdoba, Argentina
 mario@oac.uncor.edu, dgl@oac.uncor.edu, hernan@oac.uncor.edu

Draft version February 1, 2008

ABSTRACT

We compute the cluster auto-correlation function $\xi_{cc}(r)$ of an X-ray flux limited sample of Abell clusters (XBACs, Ebeling et al. 1996). For the total XBACs sample we find a power-law fit $\xi_{cc} = (r/r_0)^\gamma$ with $r_0 = 21.1 \text{ Mpc h}^{-1}$ and $\gamma = -1.9$ consistent with the results of $R \geq 1$ Abell clusters. We also analyze $\xi_{cc}(r)$ for subsamples defined by different X-ray luminosity thresholds where we find a weak tendency of larger values of r_0 with increasing X-ray luminosity although with a low statistical significance. In the different subsamples analyzed we find $21 < r_0 < 35 \text{ Mpc h}^{-1}$ and $-1.9 < \gamma < -1.6$. Our analysis suggests that cluster X-ray luminosities may be used for a reliable confrontation of cluster spatial distribution properties in models and observations.

Subject headings: cosmology-clusters-correlation function

1. INTRODUCTION

Different authors have analyzed the cluster-cluster spatial two-point correlation function finding power-law fits of the form $\xi_{cc}(r) = (r/r_0)^\gamma$ with $\gamma \simeq -1.8$ (Bahcall & Soneira 1983, Peacock & West 1992). The value of the cluster-cluster correlation length r_0 is controversial, as also is its dependence on cluster mass. This has been achieved by studying samples selected by cluster richness and the associated mean inter-cluster separation $d_c = n_c^{-1/3}$ where n_c is the mean number density of clusters. Bahcall & West 1992 and Bahcall & Cen 1992 argue for a universal scaling relation for the two-point correlation function of rich clusters where the cluster correlation length satisfies $r_0 = 0.4d_c$. At low values of $d_c \simeq 30 - 50 h^{-1} \text{ Mpc}$ the APM Cluster Survey (Dalton et al. 1994), the Edinburgh/Durham Cluster Catalog Lumsden et al. 1992 (hereafter EDCC), and the Abell 1958 and Abell, Corwin & Olowin 1989 cluster samples give similar results consistent with $r_0 = 15 - 20 h^{-1} \text{ Mpc}$. At larger d_c , however, the analyses rely only on the Abell catalog and on a cluster sample selected from the APM Galaxy Survey (Croft et al. 1997). The results of this high richness APM cluster sample are not consistent with the universal scaling relation derived from Abell clusters by Bahcall & West 1992 since only a weak dependence of r_0 on d_c is found in the rich APM cluster sample. A partial explanation for the different results between rich Abell and APM clusters could rely on the fact that the Abell catalog is subject to visual artificial inhomogeneities in contrast with the automated and well controlled APM cluster catalog (Croft et al. 1997). It should be noted, however, that given the steeper correlations for richer clusters these results are not inconsistent with the universal relation in terms of correlation amplitude.

The problems of projection effects in cluster selection (see van Haarlem et al. 1997) may be strongly overcome by selecting clusters in the X-ray rather than the opti-

cal. Moreover, given the good correlation between X-ray luminosity and cluster mass ($L_X \propto M^{4/3}$) found in both analytical solutions (Bertschinger 1985) and in numerical simulations (Navarro et al. 1995), an X-ray selected sample is suitable to study the dependence of cluster spatial correlations on mass. In this work we explore the values of r_0 in subsamples taken from the X-ray brightest Abell-type clusters of galaxies, hereafter XBACs (Ebeling et al. 1996). This sample of clusters is complete in X-ray flux, and we have selected subsets with different cuts in X-ray luminosity L_x .

2. DATA AND ANALYSIS

The X-ray-brightest Abell-type clusters of galaxies survey (hereafter XBACs, Ebeling et al. 1996) comprises 277 objects and is a 95 % complete flux limited sample. We have restricted this catalog to galactic latitudes $|b| > 25^\circ$ and X-ray flux $f > f_{cut} = 5 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the 0.1 – 2.4 keV band comprising a final sample of 248 clusters. This sample although optically selected is confirmed by the X-ray emission of the intra-cluster gas thus excluding spurious Abell clusters generated by projection effects. Also, as discussed by Ebeling et al. 1996, the XBACs sample is unaffected to first order by the incompleteness in volume of the Abell catalog at large distances since missing Abell clusters of low richness would not be included in XBACs due to their low X-ray luminosity.

In figure 1 we plot the X-ray luminosity of the clusters L_x as a function of redshift z taken from Table 3 of Ebeling et al. 1996. The smooth curve corresponds to the luminosity of an object with flux f_{cut} at redshift z in a flat cosmology. Cluster distances d were derived using the standard relation (e.g., Sandage 1961)

$$d = \frac{c(q_0 z + (q_0 - 1)((1 + 2q_0 z)^{1/2} - 1))}{h_0 q_0 (1 + z)^2} \quad (1)$$

where z is the cluster redshift, h_0 is the Hubble constant

¹CONICET, Argentina

²present address, Physics Department, University of Durham, U.K.

in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$, and c is the speed of light. Throughout this paper we have adopted a deceleration parameter $q_0 = 0.5$.




Figure 1. X-ray luminosity L_x vs redshift z for the total sample of 248 clusters analyzed. The smooth curve displays the luminosity corresponding to flux f_{cut} at redshift z in a flat cosmology.

We calculate the cluster-cluster two-point spatial correlation function $\xi_{cc}(r)$ cross correlating the data with a random catalog constructed by randomizing the angular positions of the clusters with the same redshift distribution. Each random catalog has n_{ran} points homogeneously distributed within the same boundary than the subsamples. To compute $\xi_{cc}(r)$ we have used the estimator

$$\xi_{cc}(r) = 2f \frac{n(r)}{n_{ran}(r)} - 1 \quad (2)$$

where $n(r)$ and $n_{ran}(r)$ are the number of cluster-cluster and cluster-random pairs separated by a distance r respectively; $f = N_{ran}/(N - 1)$ where N and N_{ran} are the total number of clusters in the observed sample and random catalog respectively.

We have considered 4 lower limits in X-ray luminosity which define volume incomplete subsamples of clusters (subsamples 1i to 4i). We have also defined other 4 subsamples by further imposing the restriction of a cut in redshift (z_{cut}) in order to build volume complete subsamples (subsamples 1c to 4c) (see tables 1 and 2). In figure 1 the horizontal lines define the lower limits of the 4 incomplete subsamples. The regions above the horizontal lines and with limiting redshift at the vertical lines define the 4 complete subsamples. The resulting number N of clusters is between 85 and 214 in the incomplete subsamples, and between 43 and 72 in the complete subsamples.

Sample	L_x $10^{44} h^{-2} \text{ erg s}^{-1}$	N	γ	r_0 $\text{Mpc } h^{-1}$
All	≥ 0.02	248	-1.92	$21.1^{+1.6}_{-2.3}$
1i	≥ 0.27	214	-1.89	$22.1^{+2.7}_{-2.9}$
2i	≥ 0.54	168	-1.80	$23.5^{+4.5}_{-4.7}$
3i	≥ 0.91	117	-1.59	$30.1^{+8.8}_{-10.9}$
4i	≥ 1.38	85	-1.75	$27.0^{+11.2}_{-16.0}$

Table 1. Incomplete Subsamples

We have fitted the correlation functions obtained with power laws of the form $\xi_{cc}(r) = (r/r_0)^\gamma$. We have estimated the best fitting parameters γ and r_0 and their associated errors using a maximum-likelihood estimator using a χ^2 -minimization procedure developed by Levenberg and Marquard, (see Press et al. 1987). This method deals with the errors in each distance bin providing a reliable set of fitting parameters to the correlation function. In our calculations we assume Poisson errors $\simeq \sqrt{n(r)}$ in each bin to estimate the uncertainty in the correlation function (see Croft et al. 1997 and references therein).

Sample	$L_x [10^{44}$ $h^{-2} \text{ erg s}^{-1}]$	z ($\leq$)	N	γ	r_0 $\text{Mpc } h^{-1}$
1c	≥ 0.27	0.071	59	-1.76	$26.4^{+6.9}_{-7.9}$
2c	≥ 0.54	0.101	72	-1.80	$24.6^{+6.5}_{-8.3}$
3c	≥ 0.91	0.130	54	-1.59	$30.1^{+8.8}_{-10.9}$
4c	≥ 1.38	0.160	43	-1.77	$34.7^{+12.3}_{-16.9}$

Table 2. Complete Subsamples




Figure 2. Cluster-cluster two-point correlation functions $\xi_{cc}(r)$ corresponding to the total sample.

In figure 2, 3a and 3b are shown $\xi_{cc}(r)$ corresponding to the total sample, the incomplete subsamples 1i-4i, and the complete subsamples 1c-4c respectively. Error bars in $\xi_{cc}(r)$ correspond to Poisson estimates of the uncertainties in the number statistics $\simeq \sqrt{n(r)}$.




Figure 3a. Cluster-cluster two-point correlation functions $\xi_{cc}(r)$ corresponding to the different incomplete subsamples analyzed.




Figure 3b. Cluster-cluster two-point correlation functions $\xi_{cc}(r)$ corresponding to the different complete subsamples analyzed.

Estimates of the uncertainties in the power-law best-fitting parameters r_0 and γ of the correlation functions may be visualized as plots of error contours $\chi^2 - \chi^2_{ML}$ in the $r_0 - \gamma$ plane. In figures 4, 5a and 5b we show the corresponding error contours of confidence (1, 2 and 3 σ level) of the total sample, incomplete subsamples 1i-4i, and complete subsamples 1c-4c.




Figure 4. Ellipses of confidence (1, 2 and 3 σ level) corresponding to the correlation functions shown in figures 2.




Figure 5a. Ellipses of confidence (1, 2 and 3 σ level) corresponding to the correlation functions shown in figures 3a.




Figure 5b. Ellipses of confidence (1, 2 and 3 σ level) corresponding to the correlation functions shown in figures 3b.

3. CONCLUSIONS

We have analyzed the two-point spatial correlation function of clusters of galaxies selected from a sample of X-ray brightest Abell-type clusters of Ebeling et al. 1996. For the total XBACs sample we find a power-law fit of the form $\xi_{cc}(r) = (r/r_0)^\gamma$ with $r_0 = 21.1^{+1.6}_{-2.3}$ Mpc h^{-1} and $\gamma = -1.92$, values consistent with those derived for $R \geq 1$ Abell clusters (see Bahcall & West 1992 and references therein).

In order to provide an insight of the dependence of the cluster spatial correlation length r_0 on mass, we have estimated auto-correlation functions for subsamples of clusters with different X-ray luminosity thresholds. We find

a weak increase of the correlation amplitude with increasing X-ray luminosity which is not statistically significant and suggests a lack of a strong dependence of r_0 on cluster mass. For instance, in our complete subsample 4i with the highest X-ray luminosity threshold $L_x > 1.38 \cdot 10^{44} h^{-2} \text{ergs}^{-1}$ we obtain the highest value of correlation length, $r_0 \simeq 34.7^{+12.3}_{-16.9}$ Mpc h^{-1} . Nevertheless, this value does not differ significantly from $r_0 \simeq 26.4^{+6.9}_{-7.9}$ Mpc h^{-1} corresponding to subsample 1c with $L_x > 0.2710^{44} h^{-2} \text{ergs}^{-1}$.

There is a well documented evidence for the dependence of the correlation length on cluster richness as indicated by the relation between r_0 and the mean inter-cluster separation d_c in Abell cluster samples. The weak dependence of r_0 with X-ray luminosity threshold as derived from our analysis is partially related to the broad relation between L_x and richness (Briel & Henry 1994). The relation between mass, richness and X-ray luminosity is uncertain and is affected by several observational biases and systematics (contamination by projection, departures from hydrostatic equilibrium, etc) as well as astrophysical issues (galaxy formation and evolution in clusters, pre-heating of the intra-cluster gas, shocks and supernova heating, etc). These effects are important for a suitable interpretation of the observations given the different mass dependence of the cluster correlation length expected in the variety of scenarios for structure formation. On the theoretical side the situation is also unclear. In hierarchical models of the CDM type the dependence of r_0 on d_c is found either very weak (Croft & Efstathiou 1994), or moderate (Bahcall & Cen 1992), discrepancies that according to Eke et al. 1996 may rely on the different cluster identification algorithms. These considerations and the results of our analysis suggest that cluster X-ray luminosities may be used for a reliable confrontation of models and observations.

This work was partially supported by the Consejo de Investigaciones Científicas y Técnicas de la República Argentina, CONICET, the Consejo de Investigaciones Científicas y Tecnológicas de la Provincia de Córdoba, CONICOR and Fundación Antorchas, Argentina.

REFERENCES

- Abell, G.O. 1958 ApJS, ApJ, 3, 211.
- Abell, G.O., Corwin, H.G. & Olowin R.P. 1989, ApJ, 407, L49.
- Bahcall, N.A. & Cen, R. 1992, ApJ, 398, L81.
- Bahcall, N.A. & Soneira, R.M. 1983, ApJ, 270, 20.
- Bahcall, N.A. & West, M. 1992, ApJ, 392, 419.
- Bertschinger, E. 1985, ApJS, 58, 39.
- Briel, U.G. & Henry, J.P. 1994, A&A, 278, 379.
- Croft, R.A.C. & Efstathiou, G. 1994, MNRAS, 267, 390.
- Croft, R.A.C. Dalton G.B., Efstathiou, G.P. 1997 MNRAS, in press.
- Dalton G.B., Efstathiou G., Maddox S.J. & Sutherland W.J. 1994, MNRAS, 269, 151.
- Ebeling, H., Voges, W., Bohringer, H., Edge, A.C., Huchra, J.P. & Briel, U. G. 1996, MNRAS, 281, 799.
- [Eke, R., Cole, S. and Frenk, C. 1996, MNRAS, 282, 263.
- Lumsden, S.L., Sutherland, W.J., Collins, C.A. & Guzzo, L. 1992, MNRAS, 258, 1.
- Navarro, J.F., Frenk, C.S. and White, S.D.M. 1995 MNRAS, 275, 720.
- Peacock, J.A. & West, M.J. 1992, MNRAS, 259, 494.
- Press, W.H., Flannery, B.P., Teukolsky, S.A., & Vetterling, W.T. 1987. Numerical Recipes. (Cambridge University Press, Melbourne).
- Sandage, A. 1961, ApJ, 133, 355.
- van Haarlem, M.P., Frenk, C.S. and White, S.D.M. 1997, MNRAS, 287, 81.